

AN ARSENIC-SPECIFIC BIOSENSOR WITH GENETICALLY ENGINEERED *SHEWANELLA*
ONEIDENSIS IN A BIOELECTROCHEMICAL SYSTEM

A Project Report
Presented to the Faculty of the Graduate School
of Cornell University
in Partial Fulfillment of the Requirements for the Degree of
Master of Engineering

by
Dylan Patrick Webster
May 2013

© 2013 Dylan Patrick Webster

ACKNOWLEDGEMENTS

I wish to thank my advisor, Dr. Largus T. Angenent, for providing the resources and encouragement necessary to turn my mere idea to reality. Additionally, I wish to thank Spencer Chen, Steven Chen, Claire Paduano, Caleb Radens and Swati Sureka for assisting with benchwork; Drs. Shivaun Archer, Matthew DeLisa, and Gene Madsen (of Cornell University) for providing strains and additional resources; Dr. Jeffrey A. Gralnick (of the Biotechnology Institute and Department of Microbiology, University of Minnesota—Twin Cities) for providing a *Shewanella oneidensis mtrB* deletion mutant; and Ali Awan, Devin Doud, Elliot Friedman, Taylor Stevenson, Michaela TerAvest, and Dr. Didi Waraho for providing helpful advice on experiments and protocols. This work was partially supported by the Oil Sands Leadership Initiative as part of the 2012 International Genetically Engineered Machines Competition and from an NSF CAREER grant (0939882) to Largus T. Angenent.

RESEARCH HIGHLIGHTS

- (1) We have developed the first bioelectrochemical system (BES) based environmental biosensor with genetically encoded specificity for a toxic metal.
- (2) This biosensor employs a genetically engineered *Shewanella oneidensis* strain, which was designed to produce more current in response to arsenic when cultured in a BES.
- (3) Our first-generation BES-based biosensor has a response time of approximately 24 h and a detection limit on the order of 100 μM arsenite.

ABSTRACT

Genetically-engineered microbial biosensors have yet to realize commercial success in environmental applications, due in part to difficulties associated with transducing and transmitting traditional bioluminescent information. The use of bioelectrochemical systems (BESs) in biosensing applications allows for a direct electronic output that can be more easily incorporated into devices for remote environmental monitoring. Herein, we describe the first BES-based biosensor with genetically encoded specificity for a toxic metal. By placing an essential element of the metal reduction (Mtr) pathway of *Shewanella oneidensis* MR-1 under the control of an arsenic-sensitive promoter, we have engineered a strain that produces increased current in response to arsenic when inoculated into an BES. When operated as a chemostat with a hydraulic retention time of 7 h, our BES-based biosensor has a response time of 24 h and a lower detection limit on the order of 100 μ M arsenite. To heighten the sensor's sensitivity to arsenic, we have begun characterizing an alternative sensing strain with improved translational efficiency. This ability to tune analyte sensitivity—along with the reliability of our continuous assay and the simplicity of the transcriptional circuit required for BES-based biosensing—suggests that similar sensing systems may be readily developed for both environmental deployment and on-line process control.

1. INTRODUCTION

Microbes are thought to be a promising platform for biosensing since their simplicity to cultivate, ability to thrive under adverse conditions, and amenability to modern recombinant DNA technologies suggest that microbial biosensors can offer inexpensive and specific measurement tools (D'Souza, 2001). Past microbial biosensors have been principally developed for environmental applications, since microbes have at their disposal the tools to sense both bioavailability and toxicity (Belkin, 2003). Many of these have relied on the analyte-induced expression of a reporter genes that produce a bioluminescent, fluorescent, or colorimetric signal (Joshi et al., 2009; Ramanathan et al., 1997; Sagi et al., 2003; Wang et al., 2012). Most notably, *Pseudomonas flourescens* HK44 is a well-characterized strain that has been deployed to monitor naphthalene and salicylate bioavailability in soils (Heitzer et al., 1992; King et al., 1990). Despite their popularity in academic literature, however, genetically engineered microbial biosensors have yet to see commercial success in their target environments (Shin, 2011).

One possible explanation for this gap between academia and industry is the difficulty of detecting the expression of common reporter genes—typically related to either luciferase, green fluorescent protein, or β -galactosidase (Lei et al., 2006)—remotely. Because remote monitoring requires neither on-site technicians nor transportation of samples to laboratories, the ability of microbial biosensors to transmit information across distances will likely be central in their penetration of the environmental monitoring market. Consequently, there has been recent interest in the development of deployable bioluminescent biosensors for long-term remote monitoring (Nivens et al., 2004). However, this technology has yet to gain traction—largely due

to the difficulty of creating a device that not only facilitates the transduction and transmission of bioluminescent information, but also provides an adequate environment to maintain engineered microbes for extended periods of time (Bjerketorp et al., 2006).

A promising alternative that may overcome the limitations of more traditional microbial biosensing methods is the use of bioelectrochemical systems (BESs), which take advantage of the ability of microbes to catalyze the reduction of an anode in an electrochemical cell (Friedman et al., 2012b). Most commonly, BES technologies are developed as microbial fuel cells, which are designed to harvest electricity from chemical energy stored in organic waste. Because direct electronic output is provided by BESs, however, this technology is also ideal for many biosensing purposes (TerAvest et al., 2011). While past BES-based biosensors have been designed to detect biological oxygen demand (Kim et al., 2003), carbon source availability (Kim et al., 1999), and general toxicity (Wang et al., 2013), only recently has it been shown that specific chemical signals may be detected by such sensors (Golitsch et al., 2013). Herein, we move beyond this recent proof of principle and describe a genetically encoded biosensor for arsenic that employs a *Shewanella oneidensis* strain growing in a continuous flow BES. The sensing strains are genomic $\Delta mtrB$ mutants, and are complemented by a plasmid-encoded copy of *mtrB* that is driven by an arsenic-inducible promoter. Because MtrB is essential for electrode reduction (Coursolle et al., 2010), current production is upregulated in response to arsenic.

2. MATERIALS AND METHODS

2.1 Strains and growth conditions

S. oneidensis MR-1 is an environmental isolate first found in Lake Oneida, New York (Myers and Nealson, 1988). We used an MR-1 *mtrB* deletion mutant ($\Delta mtrB$) as a basis for the construction of our arsenic-sensing strains (Coursolle and Gralnick, 2012). During experiments, we employed minimal M4 media, which contains—per liter of deionized water—0.221 g K_2HPO_4 , 0.099 g KH_2PO_4 , 0.168 g $NaHCO_3$, 1.189 g $(NH_4)_2SO_4$, 7.305 g $NaCl$, 1.192 g HEPES; 0.0713 g $CaCl_2$, and 10 mL M4 trace mineral solution. Per liter deionized water, M4 trace mineral solution contains 2.26 g Na_2EDTA , 24.89 g $MgSO_4 \cdot 7H_2O$, 0.029 g $MnSO_4 \cdot 4H_2O$, 0.058 g $NaCl$, 0.068 g $FeCl_2$, 0.065 g $CoCl_2$, 0.029 g $ZnSO_4 \cdot 7H_2O$, 0.005 g $CuSO_4 \cdot 5H_2O$, 0.35 g H_3BO_3 , 0.08 g Na_2MoO_4 , 0.119 g $NiCl_2 \cdot 6H_2O$, and 0.028 g Na_2SeO_4 (Rosenbaum et al., 2010). When appropriate, sodium L-lactate and Fe(III) citrate were added to minimal M4 media at final concentrations of 20 and 10 mM, respectively, to provide a carbon source and electron acceptor.

Escherichia coli DH5 α and WM3064 (a diaminopimelic acid (DAP) auxotroph derived from strain β 2155; Dehio and Meyer, 1997) were used for routine cloning and conjugation into *Shewanella* strains, respectively. For growth of *E. coli* WM3064, we supplemented all media with 300 μ M DAP. All *Escherichia* strains were grown at 37 °C unless otherwise noted; all *Shewanella* strains were grown at either room temperature or 30 °C. All strains were maintained on solid lysogeny broth (LB) medium at 1.5% agar. Overnight cultures were grown in 5 mL volumes of LB—either from single colonies on agar plates or from frozen stocks maintained in glycerol. When appropriate, we added kanamycin at a final concentration of 50 μ g/mL to growth media.

2.2 Mutant construction

We used standard molecular biology techniques to construct plasmids containing arsenic-responsive genetic circuits. We obtained BBa_K098994 and BBa_J33201—derived from the genomes of *S. oneidensis* and *E. coli* JM109 (Diorio et al., 1995), respectively—from the Registry of Standard Biological Parts (<http://partsregistry.org>). BBa_J33201, which contains both an arsenic-inducible promoter (P_{ars}) and its downstream transcriptional regulator (*arsR*), was amplified *via* PCR using a forward primer (ArsR_1) upstream of an EcoRI/XbaI multiple cloning site (MCS), and a reverse primer (ArsR_2) with a 5' extension introducing AscI, SpeI, and PstI cutsites (see Table S1 in Supporting Information for a list of all primers used in this study).

To facilitate the construction of the initial sensing plasmid, the coding region of *mtrB* was amplified from BBa_K098994 using a forward primer (MtrB_1) introducing AscI and BamHI cutsites flanking a ribosomal binding site (RBS) and a reverse primer (MtrB_2) introducing KpnI, SpeI, and PstI cutsites. The AscI and BamHI cutsites were included to expedite possible modifications to the RBS associated with *mtrB*. Upon digesting the resulting PCR products with EcoRI/AscI and AscI/PstI, plasmid pArsR/MtrB_1 was constructed upon the simultaneous ligation of the two fragments into pBBRBB (Addgene plasmid 32549; Vick et al., 2011), a broad host range mobile vector conferring kanamycin resistance, derived from pBBR1MCS-2 (Kovach et al., 1995). A second plasmid (pArsR/MtrB_2) was also constructed—without a BamHI cut site between the RBS and start codon associated with *mtrB*—*via* the direct digestion of the *mtrB* coding sequence from BBa_K098994. This plasmid was constructed to control for possible changes to translational efficiency with the modified *mtrB* upstream sequence in pArsR/MtrB_1.

Plasmids pArsR/MtrB_1 and pArsR/MtrB_2 were electroporated into electrocompetent *E. coli* WM3064 stocks—prepared by employing a protocol described by Wu et al. (2010)—in 1 mm cuvettes at a voltage of 1.8 kV, a capacitance of 25 μ F, and a resistance of 200 Ω . After selecting for successful transformants, the resulting *E. coli* strains were grown overnight and plated on LB+DAP in a suspension with *S. oneidensis* Δ mtrB—after being washed of residual kanamycin. Each conjugation was allowed to proceed for 8 h at 30 °C, after which a new LB+kanamycin plate, without DAP, was streaked for single colonies of *S. oneidensis*. Successfully conjugated *S. oneidensis* colonies were confirmed *via* colony PCR and Sanger sequencing.

2.3 Iron(III) citrate reduction

For high-throughput characterization of our arsenic sensing strains, we quantified their capacities to reduce Fe(III) at concentrations of arsenite ranging from 0 to 500 μ M. While the pathways responsible for electrode reduction and soluble iron reduction do not fully overlap, MtrB plays a crucial role in both pathways, making the quantification of iron reduction ideal for our purposes (Rosenbaum et al., 2010; Coursolle and Gralnick, 2012). In preparation for the assay, all strains were cultured aerobically overnight in LB at an arsenite concentration to be employed in the subsequent assay. After growing to an OD₆₀₀ of 1, cultures were washed and resuspended to a final OD₆₀₀ of 0.015 in minimal M4 media with 20 mM sodium lactate and 10 mM Fe(III) citrate. Inside an anaerobic chamber, 30 μ L of each cell suspension was added to 270 μ L of minimal M4 media (and arsenite, where appropriate) in the wells of sterile 96-well plates. After incubating for 12 h, 5 μ L from each well were transferred into clean wells of a new 96-well

plate, along with 45 μ L of 0.5 M HCl and 300 μ L ferrozine reagent (Coursolle et al. 2010, Stookey, 1970). For quantification, the resulting mixture was measured spectrophotometrically at 562 nm. Defined Fe(II) standards at an appropriate range of concentrations were used to calibrate the assay.

2.4 BES construction and electrochemical techniques

We employed single-chambered glass electrochemical reactors in the construction and characterization of our bench-scale BES-based biosensor. Operating BESs as electrochemical half-cells, we employed either a commercial potentiostat (VSP, Biologic USA, Knoxville, TN; CHI660B, CH Instruments, Austin, TX) or an open-source, microcontroller-based potentiostat (placed in a Faraday cage for electronic shielding; Friedman et al., 2012a) to control and monitor electrochemical conditions in a three electrode system. We controlled the potential of a graphite working electrode (Poco Graphite, Inc., Decatur, TX) at 0.3 V *versus* a Ag/AgCl, saturated KCl glass reference electrode, while a graphite counter electrode served as a current drain. In this arrangement, *S. oneidensis* biofilm formation and electron transfer is induced at the working electrode, since it is poised at a potential that is energetically favorable for electrode reduction.

In continuous flow experiments, the working volume was held constant at 120 mL while media was fed *via* a peristaltic pump (Cole-Parmer, Vernon Hills, IL) at a continuous rate of 18 mL/h to maintain a hydraulic retention time of about 7 h. Because recent evidence suggests that oxygen intrusion allows *S. oneidensis* to overcome redox mediator washout in continuous-flow reactors (TerAvest, unpublished data), the head space of the reactor was left open to atmosphere

through a 0.2 μm filter (Pall Corporation, Cortland, NY) to maintain a micro-aerobic environment. This environment also provided the sensing strains with oxygen as an alternative to the electrode as an electron acceptor, so that electron acceptors would not be limiting when arsenic was not present (*i.e.*, in conditions where the Mtr pathway was incomplete).

We operated reactors for continuous-flow investigations at an initial arsenite concentration of zero. Current was recorded *versus* time for BESs inoculated with a *S. oneidensis* strain carrying pArsR/MtrB_1, as well as with ΔmtrB and MR-1 empty vector controls. We added kanamycin to the reactor feed to maintain a selective pressure for plasmid maintenance. After observing stationary current (with fluctuations within 5% of average value) for more than two hydraulic retention times, arsenite was added directly to a feed tank such that the steady-state arsenite concentration inside the reactors could be controlled. These steady-state arsenite concentrations were incrementally increased (again upon observation of stationary current) until a final concentration of 500 μM was reached.

We also performed batch investigations of a *S. oneidensis* strain carrying pArsR/MtrB_2, alongside as ΔmtrB and MR-1 empty vector controls. *S. oneidensis* strains were grown overnight in LB+kanamycin (without arsenite) to an OD_{600} just under 1.0 before being resuspended in minimal M4 (with kanamycin) to an OD_{600} of 0.015. Suspensions were inoculated into 100 μL of minimal M4 media with kanamycin and arsenite (when appropriate) after at least 5 h of abiotic background current was observed. While current was recorded over the entire length of batch experiments, peak current was of particular interest as a response variable to arsenite concentration.

3. RESULTS AND DISCUSSION

3.1 Design of an arsenic-sensing genetic circuit

We relied on an *mtrB* complementation strategy to construct arsenic-responsive genetic circuits in plasmids pArsR/MtrB_1 and pArsR/MtrB_2. MtrB plays an essential role in the metal reduction (Mtr) pathway of *Shewanella* spp., stabilizing a complex formed between MtrA and MtrC (Coursolle et al., 2010). Because this stable complex is required for electrode reduction, strains deficient in the *mtrB* coding sequence are unable to produce significant levels of current when inoculated into BESs. However, when *mtrB* is re-introduced into a knockout strain (and transcription is induced), the electrode reduction phenotype is restored—*i.e.*, when inoculated into an BES, current production will increase in response to increasing *mtrB* transcription in such an engineered strain. We exploited this by placing the *mtrB* coding sequence under the control of an arsenic-inducible promoter region (the molecular details of which are discussed in Section 2.2).

The arsenic-inducible promoter (P_{ars}) is negatively regulated by ArsR (Fig. 1). When arsenic is excluded from a cell with these genetic components, the binding kinetics between ArsR and its associated operator region within P_{ars} are relatively strong, blocking the transcription of downstream genes (*i.e.*, *arsR* and *mtrB*). In this arrangement, ArsR acts as a negative auto-regulator by limiting its own expression to a low, basal level. The dynamics of the circuit change, however, when arsenic (in the form of arsenates or arsenites) enters the cell. In this condition, arsenic associates with ArsR, inducing a conformational change that leads to its dissociation from the P_{ars} operator. Because the binding kinetics between ArsR and the P_{ars}

operator become more unfavorable (to a point) with increasing arsenic concentrations, MtrB levels—and hence, electrode reduction capacities—increase with increasing arsenic presence.




Fig. 1. Design of arsenic-responsive transcriptional circuit in plasmids pArsR/MtrB_1 and pArsR/MtrB_2. *arsR* is a negative auto-regulator, inhibiting the transcription of itself and downstream *mtrB*. (A) When relatively little arsenic is present, ArsR is able to bind to the operator of P_{ars} , the arsenic-sensitive promoter. (B) As more arsenic enters the cell, ArsR dissociates from the operator, leading to an increase in *mtrB* expression.

3.2 Effect of arsenic on iron reduction

S. oneidensis strains carrying arsenic-responsive plasmids had increasing capacity to reduce Fe(III) to Fe(II) at increasing arsenite concentrations (Fig. 2). While a statistically significant change in iron reduction capacity was only observed at concentrations above 100 μ M for the

strain carrying pArsR/MtrB_1, the strain carrying pArsR/MtrB_2 was able to reduce significantly more Fe(III) at arsenite levels below 80 μM —compared to basal iron reduction. This indicates that the efficiency of MtrB translation plays a role in determining arsenic sensitivity, as further described in Section 3.4. However, because *both* arsenic-sensing strains were able to reduce more iron at increasing arsenite concentrations, we have strong evidence that each strain can function as the biological component of an arsenic sensor with direct electronic output when inoculated into an BES.




Fig. 2. Fe(II) evolution by *S. oneidensis* strains over 12 h at various concentrations of arsenite. (A) A significant improvement in iron reduction is observed for a *S. oneidensis* strain carrying pArsR/MtrB_1 at arsenite concentrations above 100 μM . (B) A wild-type positive control (blue) and a $\Delta mtrB$ negative control (orange) each show toxic responses to arsenic, as iron reduction capacity is diminished with increasing arsenite concentration; a *S. oneidensis* strain carrying pArsR/MtrB_2 (green) shows an increasing capacity to reduce iron at increasing arsenite concentrations, with significantly different Fe(II) evolution at 80 μM arsenite.

It is interesting to note that increasing MtrB expression can compensate for arsenic toxicity in our sensing strains—in terms of respiration rates. Both wild-type *S. oneidensis* MR-1 and the $\Delta mtrB$ negative control showed decreasing iron reduction rates at increasing arsenite concentrations (Fig. 2B), indicating that toxic effects of arsenite cause a growth defect in *S. oneidensis* strains at the range of concentrations under investigation. Therefore, it is likely that increasing MtrB expression “makes up” for this defect in our sensing strains. At arsenite concentrations above 100 μ M, however, the *S. oneidensis* strain carrying plasmid pArsR/MtrB_2 shows diminished capacity to reduce iron (data not shown). This phenomenon may be explained by saturation in MtrB activity (disallowing compensation for arsenic toxicity) combined with the inherent toxic effects of overexpression of Mtr proteins—as recently investigated by Goldbeck et al. (2013).

3.3 Effect of arsenic on continuous flow BES current production

When inoculated into a continuous flow BES, the *S. oneidensis* strain carrying plasmid pArsR/MtrB_1 reproducibly generated increased steady-state current in response to increasing steady-state arsenite concentrations. An arsenite concentration of 500 μ M induced a steady-state current of approximately 150% of the basal level (*i.e.*, current produced in an BES without arsenic; Fig. 3). While arsenite concentrations on the order of 100 μ M induced a transient increase in current, an elevated steady-state current level was not achieved. While a definitive explanation for this phenomenon has yet to be tested experimentally, we hypothesize that changing dynamics in the menaquinone pool (which supplies the Mtr pathway




Fig. 3. Response of *S. oneidensis* strain carrying pArsR/MtrB_1 to arsenite in two replicates. After panning through a steady-state arsenite concentration of 100 μM , a final response was observed at 500 μM arsenite. After a transient phase on the order of a day, a new steady-state—about 50% above baseline current—was achieved.

with reducing equivalents to be shuttled outside of the cell as current) accounts for this transient behavior. Despite these complications, we have shown that steady-state current may be recorded as an indicator of arsenite concentration, forming the basis of our BES-based biosensor.

However, limitations of these sensing capabilities include a relatively long response time (on the order of 1 day). While it is possible that the biofilm turnover rate limits the ability of our biosensor to respond rapidly to changing arsenite concentrations, some factors contributing to response time are unknown. Because the engineered *S. oneidensis* strain is deficient in MtrB at low levels of arsenic, it is also possible that changes in transcriptional regulation (in multiple stages) are required before a fully functional Mtr pathway is able to form, further contributing to the time required to reach a new steady-state upon arsenic addition. Further investigation is required, however, to confirm the role of transcriptional regulation in the response time of the

biosensor. Such investigation, along with variation of the macro-scale parameters of the reactor (e.g., biofilm shear stress due to the flow of media) will lead to improved response time in future BES-based biosensors relying on engineered *S. oneidensis* strains.

3.4 Electrochemical characterization of alternative sensing strain

In addition to the relatively long response time of the continuous flow BES-based biosensor, the detection limit on the order of hundreds of μM arsenite is also of concern. For practical applications, such a biosensor would need to be much more sensitive, because both the USEPA and WHO have set standards for maximum chronic exposure to arsenic salts that are orders of magnitude below the detection limit of the *S. oneidensis* strain carrying plasmid pArsR/MtrB_1 (McGuigan et al., 2010). Consequently, the arsenic-responsive circuit in pArsR/MtrB_1 requires modification to increase arsenic sensitivity.

As mentioned in Section 2.2, the *mtrB* upstream sequence was modified in the insertion of a BamHI cut site in pArsR/MtrB_1. The nucleotide changes in this altered upstream sequence are shown in Table 1. Because KineFold (<http://kinfold.curie.fr/>) simulations predict

	<i>mtrB</i> Upstream Sequence
Native	AGGAGA AATACTAGAT <i>G</i>
Altered	A AGGAGA <u>CGGATCC</u> AT <i>G</i>

Table 1. *mtrB* upstream sequences for native (wild type and pArsR/MtrB_2) and altered (pArsR/MtrB_1) genotypes. During the construction of pArsR/MtrB_1, the number and identity of spacer nucleotides flanked by the ribosomal binding site (RBS; bold) and the start codon (italic) were changed upon introduction of a BamHI cut site (underlined).




Fig. 4. Current production over time for *S. oneidensis* strain carrying plasmid pArsR/MtrB_1 in batch BESs. Three replicates are shown in each panel. When inoculated into a reactor without arsenic (A), maximum current production is consistently below 20 μA for this engineered strain; in reactors with 40 μM arsenite (B), maximum current is around 20 μA ; in reactors with 100 μM arsenite (C), maximum current is above 30 μA .

mRNA folding to partially block access to the RBS of *mtrB* for this sequence, it is likely that strains carrying pArsR/MtrB_1 exhibit reduced MtrB translational efficiency. As mentioned in Section 2.2, however, pArsR/MtrB_2 retained the native *mtrB* upstream sequence; strains carrying this plasmid should not exhibit such reduction. This was partially confirmed by the results of iron reduction assays presented in Section 3.2, since strains carrying pArsR/MtrB_2 displayed a significantly higher capacity to reduce iron at relatively low arsenite concentrations—

suggesting that higher MtrB concentrations were achieved. Further evidence for the enhancement in translational efficiency conferred by pArsR/MtrB_2 is shown in Fig. 4 above, which displays current production in batch BESs at three arsenite concentrations.

While further controls and replicates are required to achieve statistically significant results, preliminary evidence makes it clear that the *S. oneidensis* strain carrying pArsR/MtrB_2 is able to respond to arsenic concentrations on the order of tens of μM , a full order of magnitude less than the detection limit of the original strain. While further work is required to improve the detection limit of our BES-based biosensor, it is encouraging that modulation of translational efficiency is an apparently effective approach in the tuning of biosensor sensitivity. Because the construction of RBS libraries (characterized in specific hosts) is already underway as part of the Registry of Standard Biological Parts, this implies that synthetic circuits with modulated sensitivity will be relatively easy to build as the burgeoning field of synthetic biology expands.

4. CONCLUSIONS

We have shown that it is possible to transduce chemical information—of relevance to environmental quality—into the digital world by employing a BES inoculated with genetically engineered *S. oneidensis*. By characterizing current responses of our engineered strains to arsenic, we have moved beyond a proof of principle, and have constructed an BES-based biosensor with direct environmental applications. Because our biosensor outputs current directly, it may be more easily integrated into a deployable device for remote monitoring than more traditional microbial biosensors. Such a prototype (designed for the remote monitoring of arsenic in natural

bodies of water) was constructed as part of the 2012 International Genetically Engineered Machine (iGEM) Competition by Cornell iGEM (<http://2012.igem.org/Team:Cornell>; Fig. 5). Briefly, this device is capable of supporting monitoring for up to six




Fig. 5. Overview of field-deployable biosensor developed for 2012 iGEM competition. An engineered, arsenic-responsive *S. oneidensis* strain is inoculated into a reactor housed in a rugged device—alongside a control strain to account for fluctuations in environmental parameters. Sterile lactate is fed to reactors from vessels built into the device, combining with a filter-sterilized sample stream. On-board electronics record current over time, and transmit data wirelessly so that arsenic concentrations can be monitored remotely.

months—maintaining a continuous culture of an arsenic-sensing *S. oneidensis* strain with a concentrated stock of substrate—and is capable of wirelessly transmitting water quality data by employing a modified version of a previously published open-source potentiostat (Friedman et al., 2012a). Because such a device is less expensive and labor intensive than more conventional analytic chemistry approaches, it is clear that BES-based biosensors are ideal for deployment as a

first-pass monitoring system for remote detection of contamination events. As Golitsch et al. (2013) note, because *S. oneidensis* strains can be modified to detect a multitude of analytes of environmental interest with the simple swap of a promoter, a number BES-based biosensors may be constructed by hijacking the regulation of the Mtr pathway of *S. oneidensis*. Because of this, along with the fact that the genetic system of this organism is well worked-out, we believe *S. oneidensis* to be an ideal host for development of biosensors for remote monitoring of environmental contaminants.

5. APPENDIX A. SUPPORTING INFORMATION

Primer	Restriction Site(s)	Sequence
MtrB_1	AscI/BamHI	TTCTTCGGCGCGCCTAAGGAGACGGATCCATGAAATTTAAACTCAATTTGATCACTC
MtrB_2	KpnI/SpeI/PstI	GTTTCTTCCTGCAGTCAGACGAACTAGTAGGTACCATAGAAGGTGAGCCAGTGTGAC
ArsR_1	EcoRI	CGTATCACGAGGCAGAATTC
ArsR_2	AscI/SpeI/PstI	GTTTCTTCCTGCAGTACTAGTAGGCGCGCCGTTAACTGCAAATGTTCTTACTGTCC

Table S1. Primers and restriction sites used for cloning and subsequent generation of arsenic-sensing strains. MtrB_1 and MtrB_2 were used to amplify *mtrB* from [BBa_K098994](#) (Registry of Standard Biological Parts); ArsR_1 and ArsR_2 were used to amplify *arsR* from [BBa_J33201](#) (Registry of Standard Biological Parts).

5. 1 GenBank Formatted Sequence for Plasmid [pArsR/MtrB 1](#):

```

LOCUS       p_arsR/mtrB_p14             6525 bp    DNA     circular UNA 20-MAR-2013
DEFINITION   Arsenic inducible expression of mtrB; Derived from pBBRBB-eGFP.
ACCESSION    urn.local...1363788224712.13
VERSION      urn.local...1363788224712.13
KEYWORDS     .
SOURCE       .
  ORGANISM   .

FEATURES             Location/Qualifiers
     misc_feature     complement(<1..305)
                       /note="Geneious type: ORF"
                       /label="Rep"
     CDS              <1..389
                       /translation="MPQQRFLVVVINHHGGTEHRQAQLVAGLAPRHVIDHVGRRHGA
                       GAVELHDGDPALGHQVLDVLDRLPQRTSDELGKCLLADHHGVLVAHLRHEVMQQHCR
                       GFPRNKGPRMLRFAFRLHPVTGLVLGLNADFSGLRGHAYLHAGVCRVTAAAPCAISCN
                       FSAARQLCVA*"
                       /label="ORF frame 3"
     CDS              complement(<1..305)
                       /translation="MEISMATQSREIGIQAKNKPGRHWVQTERKAHEAWAGLIARKPTA
                       AMLLHLHVAQMGHQNNAVVSQKTLKSLIGRSLRTVQYAVKDLVAERWISVVKLNGPGT
                       VSAVVVNDRVAGWQPRDQLRLSVFSAAVVVDHDDQDESLLGHGDLRRIPTLYPGEQQL
                       PTGPGEEPSPQPGIPGMEFDLPALTETEWEERRGQRLPMPDEPCFLDDGEPELEPPT
                       VTLPBR*"
                       /label="ORF frame 2"
     misc_feature     894..2285
                       /note="Geneious type: ORF"
                       /label="mob"
     CDS              894..2285
                       /translation="MGQGRPRPSWSRWVCGRRLLGSRWAIPVVKLVGRRSLISLSSRV
                       VHGPEASQPVAAARGHRPHIHGLARERSDRAGRSPESKPVGRRSRRRSLCEAKSSE
                       YTQALSGPPEAPCAAPVEPVGHQKGAGMAAYAIMRCKKLAKMGNVAASLKHAYRER
                       ETPNADASRTPENEHMAASSTDEAMGRLELLPEKRRKDAVLAVEYVMTASPEWWSA
                       SQEQQAFFFEKAHKWLADKYGADRIVTASIHRDETSQHMTAFVVPVLTQDGRLSAKEFI
                       GNKAQMTDRDQTTFAAAVADLGLQRIEGSKARHTRIQAIFYALERPPVGHVTISPQAV
                       EPRAYAPQGLAEKLGISKRVETPEAVADRLTKAVRQGYEPALQAAAGAREMRKKADQA
                       QETARDLRERLKPVLDALGPLNRDMQAKAAAIKAVGEKLLTEQREVQRQKQARQQE
                       RGRAHFPEKCHLG*"
                       /label="ORF frame 3"
     promoter         2405..2454
                       /label="NEOKAN_promoter"
     CDS              2543..3337
                       /translation="MIEQDGLHAGSPAAWVERLFGYDWAQQTIGCSDAAVFRLSAQGR
                       PVLVKTDLGALNELQDEAARLSWLATGVPCAAVLDVVTEAGRDWLLGEVPGQDL
                       LSSHLPAAEKVSIADAMRRLHLDPATCPFDHQAHRIERARTRMEAGLVDQDDLE
                       EHOGLAPEALFARLKARMPDGEDLVVTHGDACLPIIMVENGFRSGFIDCGRLGVADRY
                       QDIALATRDIAEELGGEWADRFLVLYGIAAPDQRIAFYRLLEFF*"
                       /label="ORF frame 2"
     gene             2546..3334
                       /gene="NeoR/KanR"
                       /label="NeoR/KanR"
     CDS              complement(2852..3388)
                       /translation="MAGWASLGRSFRTPESRSEELVKKAIEGDALRIGSGDTVKHEEA
                       VSPFLAKLFSNITGSQRYVLIARHTQPATVDESRAAIFHHDIRQAGIAMGHDEILA
                       VGHARLEPGEQFGWREPLMLFVQIILIDKTGFHPSTCSLDAMFRLVVEWAGSRIKRMQ
                       PPHCISHDGYFLGRSKVR*"
                       /label="ORF frame 3"
     misc_feature     3499..3529
                       /note="Geneious type: operator"
                       /label="ArsR binding site"
     RBS              3566..3571
                       /label="RBS"
     gene             3577..3930
                       /label="arsR"
     primer_bind      3955..3982
                       /label="FOR_PRIME_mtrB_RBSswap_V2"
     gene             3955..6048
                       /label="mtrB"

ORIGIN
1  gtgccggggc  cgttgagctt  caccacggag  atccagcgct  cggccaccaa  gtccttgact
61  gcgtattgga  ccgtccgcaa  agaactgccg  atgagcttgg  aaagtgtctt  ctggctgacc
121  accacggcgt  tctggtggcc  catctgcgcc  acgaggtgat  gcagcagcat  tgccgccgtg
181  ggtttctctg  caataagccc  ggcccacgcc  tcatgcgctt  tgcgttccgt  ttgcacccag
241  tgacccggct  tgttcttgcc  ttgaatgccg  atttctctgg  actgcgtggc  catgcttato
301  tccatgcggt  aggggtccgc  acggttgccg  caccatgcgc  aatcagctgc  aacttttcgg
361  cagcgcgaca  acaattatgc  gttgcgtaaa  agtggcagtc  aattacagat  tttctttaac
421  ctacgcaatg  agctatttgc  gggggtgccg  caatgagctg  ttgcgtacc  ccccttttta
481  agttgttgat  ttttaagtct  ttgcgatttc  gccctatatc  tagttctttg  gtgcccacaa
541  aaggggcacc  ctgcgggggt  ccccccacgc  ttccggcgcg  ctccccctcc  ggcaaaaaat
601  ggcccccctc  gggcttggtg  atcgactcgc  cggccttcgc  ccttgcccaa  ggtggcgctg
661  cccctctgga  acccccgcac  tcgcccgctg  gaggtctcgg  gggcaggcgg  gcgggcttcg
721  ccttcgactg  ccccactcgc  cataggtctg  ggtcgttcca  ggcgcgtcaa  ggccaagccg
781  ctgcgcggtc  gctgcgcgag  cettgacccg  ccttccactt  ggtgtccaac  cgccaagcga
841  agcgcgcgag  ccgcaggccg  gaggtctttc  cccagagaaa  attaaaaaaa  ttgatggggc
901  aaggccgcag  gccgcgcagt  ttgagccggt  gggatatgtg  tcgaaggctg  ggtagccggt
961  gggcgaatcc  tgtgttcaag  ctctgtggga  ggccgagcct  gtccatcagc  ttgtccagca
1021  ggggttgtcc  cgggcccagc  gaagcgagcc  agccggtggc  cgctcgcggc  catcgtccac
1081  atatccacag  gctggcaagg  gagcgagcgc  acccgcgagg  gcgaagccgc  gagagcaagg

```

1141 ccgtagggcgc ccgcagccgc cgtagggcgt caccactttg cgaagcaaaag tctagtgagt
1201 atactcaagc attgagtggc ccgcgcggag caccgccttg cgtcgcccc gtcgagccgg
1261 ttggacacca aaagggaggg gcaggcatgg cggcatacgc gatcatgcga tgcagaagc
1321 tggcgaaaat gggcaacgtg gcggccagtc tcaagcacgc ctaccgcgag cgcgagacgc
1381 ccaacgctga ccgcagcagg acgccagaga acgagcactg ggcggccagc agcaccgatg
1441 aagcgatggg ccgactgcgc gacttgctgc cagagaagcg gcgcaaggac gctgtgttgg
1501 cggtcgagta cgtcatgacg gccagcccg aatggtggaa gtcggccagc caagaacagc
1561 aggcggcggt ctccgagaag gcgcacaagt ggctggcgga caagtacggg gcggatcgca
1621 tcgtgacggc cagcatccac cgtgacgaaa ccagcccgcga catgaccgcg ttcgtggtgc
1681 cgctgacgca ggacggcagg ctgtcgccca aggagttcat cggcaacaaa gcgcagatga
1741 cccgcgacca gaccacgttt gcggctgctg tggccgatct agggctgcaa cggggcgcag
1801 agggcagcaa ggcacgtcac acgcgcattc aggcgttcta cgaagccctg gagcgccac
1861 cagtgggcca cgtcaccatc agcccgcagg cggctcgagcc acgcgcctat gcaccgcagg
1921 gattggccga aaagctggga atctcaaaag cgttgagac gccggaagcc gtggccgacc
1981 ggctgacaaa agcggttcgg caggggtatg agcctgccct acaggccgcc gcaggagcgc
2041 gtgagatggc caagaagccc gatcaagccc aagagacggc ccgagacctt cgggagcgc
2101 tgaagcccggt tctggaagcc ctggggcgct tgaatcggga tatgcaggcc aaggccgcg
2161 cgatcatcaa ggcctggggc gaaaagctgc tgacggaaca gcgggaagtc cgcgcgaga
2221 aacagggcca gcgccagcag gaacgcgggc gcgcacattt ccccgaaaag tgccacctgg
2281 gatgaatgct agctactggg ctactctggac aaggggaaaac gcaagcgcaa agagaaagca
2341 ggtagcttgc agtgggctta catggcgata gctagactgg cgggttttat ggacagcaag
2401 cgaacgggaa ttgccagctg gggcgccctc tggtaagggtt ggggaagccct gcaaaagtaa
2461 ctggatggct ttcttggcgc caaggatctg atggcgagc ggatcaaat ttgatcaaga
2521 gacaggatga ggatcgttct gcatgattga acaagatgga ttgcacgcag gttctccg
2581 cgcttgggtg gaagggctat tcggctatga ctgggcacaa cagacaatcg gctgctctga
2641 tggcccgctg ttccggctgt cagcgagggg gcgcccgggt cttttgtca agaccgacct
2701 gtccggtgct ctgaatgaac tacaggacga ggcagcgcg cttactgtgg tgggccagc
2761 gggcgttctc tgcgcagctg tgctcgagtc tgctactgaa gcgggaaggg actggctgct
2821 attggcgcaa gtgcgggggc aggatctctc gtcatctcac ctgtctcctg ccgagaaagt
2881 atccatcatg gctgatgcaa tgccggcggt gcatacgctt gatccggcta gctgccctat
2941 cgaccaccaa gcgaacatc gcatcgagcg agcacgtact cggatggaa cgggtcttgt
3001 cgatcaggat gatctggacg aagagcatca ggggctcgcg ccagccgaac ttgttcgcca
3061 gctcaagcgc cgcattcccc acggcgagga tctcgtcgtg acccatggcg atgctgctt
3121 gccgaatatc atggtggaaa atggccgctt ttctggattc atcgactgtg gccggctggg
3181 tggcgcgac ccgtatcagg acatagcgtt ggctacccgt gatattgctg aagagcttgg
3241 cggcgaatgg gctgaccgct tctcgtgctt ttacggtatc gccgctcccc attcgacgc
3301 catcgccctc tatcgccctc ttgacgagtt cttctgagcg ggaactctgg gttcgaaagt
3361 accgaccaag ccagcccaac cctgcatca ccagatttgc attccaccgc cgccttctat
3421 gaaaggttgc actagagaat tcgcggcgcc ttctagagcc aactcaaaat tcaacatcat
3481 taccctcttc tgcaactaca cattcgttaa gtcatatatg tttttgactt atccgcttgc
3541 aagagagaca ctacctgcaa caatcaggag cgcgaatatg catttctgtt acccatccaa
3601 ttgttcaaaa ttcttgctga tgaacccgtt ctgggcatcg ttttactgct cagcgaactg
3661 ggagagttat gcgtctgcga tctctgcaat gctctcgacc agtcgacgcc caagatctcc
3721 gcgcacgtgg cattgctgcg tgaagcggg ctattgctgg accgcaagca aggttaagtgg
3781 gttcattacc gcttatcacc gcatattcca gcatggcgcg cgaataattat tgatgagggc
3841 tggcgatgtg aacaggaaaa ggttcaggcg attgtcgca acctggctcg acaaaactgt
3901 tccggggaca gtaagaacat ttgcagttaa cggcgccgct aaggagacgc atccatgaaa
3961 tttaacctca attgatcac tctagcgtta ttgccaaca caggcttggc cgtcgctgct
4021 gatggttatg gtctagcgaa tgccaatact gaaaaagtga aattatccgc atggagctgt
4081 aaagcgtcgc tcgttgaaac gggcacatca ggcactgtgg gtgtcggtgt cgggtataac
4141 agcgaagagg atattcgctc tgccaatgco tttggtacat ccaatgaagt ggcgggtaaa
4201 ttgatgcgc atttaaacct taaaggtgaa aagggttatc gtgcagatgt tgatgcttat
4261 caactcggtg ttgatggcgc tcgcttagat gtcaatgcgg gcaaacaaag ccagtacaac
4321 gtcaatgtga actatcgcca aattgctacc tacgacagca atagcgccct atcgccctac
4381 gcgggttatg gtggcaataa cctcacgtta ccggataact ggataacagc aggttcaagc
4441 aaccaaatgc cactcttgat ggacagcctc aatgcctcg aactctact taaactgtgag
4501 cgcacggggt tgggatttga atatcaaggt gaatccctgt ggagaccta tggttaactac
4561 atgcgtgaag agaaaacggg cttaaaaaaa gcctctggta gctcttcaaa ccaatcgatg
4621 atgttagcag agccggtgga ttacaccact gacaccattg aagcgggtgt caaactcaag
4681 ggtagctgtt gggttacccg actcagttac aatgggtcaa tattcaaaaa cgaatacaac
4741 caattggact ttgaaaatgc ttttaacccc acccttgggt ctcaaaacca aggtacgatg
4801 gcactcgact cggataacca gtcacacacc gtgtcgctga tgggacagta caacgatg
4861 agcaacgcac tgtcgggtgc tattctgacc ggacaaatga gccaaatga ggcgttagtg
4921 acggataact accggttatg taatcagctc aataccgatg ccgtcgatgc caaagtcgat
4981 ctactgggta tgaacctgaa agtcgttagc aaagtgaagc atgatcttgc cttaacaggt
5041 agttacgatt attacgacgc tgacaataat acccaagtag aagaatggac tcagatcagc
5101 atcaacaatg tcaacggtaa ggtggcttat aacacccctt acgataatcg tacgcaacgc
5161 tttaaaagtg ccgcagatta tcgcattacc cgcgatatac aactcgatgg ttggttatgac
5221 ttcaaacgtg accaacgtga ttatcaagac cgtgaaacca cggatgaaaa taccgtttgg
5281 gcccgtttac gtgtaaacag cttcgatact tgggacatgt gggtaaaagg cagttacggt
5341 aaccgtgagc gtcacaata ccaagcgtct gaatggacct cttctgaaac caacagcgtg
5401 ttacgtaagt acaatctggc tgaccgtgac agaactcaag tcgaagcacg gatcacccat
5461 tcgcatttag aaagcctgac tatcgatgtt ggtgccggtt acgctttaga tgattatacc
5521 gatactgtga ttggtataac tgagtcaaaa gacaccagtt atgatgcaa catcagttat
5581 atgatcacgc ctgacttact ggcaaccgac ttctacaatt accaaacctat tgagtctgaa
5641 caggcggtga gcagcaatta cagcaaccoca acgtggacag gctttataga agatcaagga
5701 gatgtgtcgc gtgcaggtat cagctacaac aatctcgttg agaacaagtt acgctagga
5761 ctggaactaca cttattccaa ctccgacagt aacactcaag tcagacaagg tatcactggc
5821 gactatgggt attattttgc caaagtgcac aacattaact tatacgtcca atatcaagcc
5881 accgagaatac tcgcgctgcg cttcgattac aaaaattgaga actataagga caatgacgcc
5941 gcaaatgata tcgccgttga tggcatttgg aacgtcgtag gttttggtag taacagccat
6001 gactcacacgc cacaatgct gatgctgagc atgagttaca aactctaata ctgagtcac
6061 actggctcac cttctatggt acctaactagt tcgtctgact gcagctagt atagtctgga
6121 acagcgcaet tacgggttgc tgcgcaacc aagtctaccc ggcgcggcag cgtgacccgt
6181 gtccggcggt ccaacggctc gccatcgctc agaaaaacag gctatcgagg catcgccagg
6241 cgtgctgccc cgcgcggttc ccattctctc gtttcgggta aggtggcagc gttctggttc
6301 atgcccggaa tgccgggctg gctggcgccg tctctgcggg ggcggctcgg tagttgctgc
6361 tcgcccggtg acagggctcg gatgcggcgc aggtcgccat gccccaacag cgtattctgc
6421 tggctcgtgt gatcaaccac cagcgcgcca ctgaacacgc acaggcgcaa ctggtcgcg
6481 ggctggcccc agccaacgcg gtcattgacc acgtaggcgc acagc

//

5.2 GenBank Formatted Sequence for Plasmid [pArsR/MtrB 1:](#)

```

LOCUS       p_arsR/mtrB (p16)             6604 bp    DNA     circular UNA 20-MAR-2013
DEFINITION   Arsenic inducible expression of mtrB; Derived from pBBRBB-eGFP.
ACCESSION    urn.local...1359230126763.14
VERSION      urn.local...1359230126763.14
KEYWORDS     .
SOURCE       .
  ORGANISM   .

FEATURES             Location/Qualifiers
     CDS             <1..389
                     /translation="MPQQRFLVVVNHGGTEHRQAQLVAGLAPRHAVIDHVGRHGA
                     GAVELHDGDPALGHQVLDCVLDPRQRTSDELGKCLADHHGVLVAHLRHEVMQQHCRR
                     GFPRNKGPRFLMRFAFRLHPVTGLVLGLNADFSGLRGHAYLHAVGCRTVAAPCAISCN
                     FSAARQQQLCVA*"
     CDS             /label="ORF frame 3"
                     complement(<1..305)
                     /translation="MEISMATQSRIGIQAKNKGHWQVTERKAHEAWAGLIARKPTA
                     AMLLHLHLVAQMGNHNAVVSQKTLKSLGRSLRTVQYAVKDLVAERWISVVKLNGPQT
                     VSAVVVNDRVAVGQPRDQLRLSVFSAVVVDHDDQDESLGHGDLRRIPTLYPGEQQL
                     PTGFGEPEPSQPGIPGMEFDLPALTETEWEERRGQQLRPFMPDEFCLDDGEFLEPPT
                     VTLPRR*"
     CDS             /label="ORF frame 2"
                     894..2285
                     /translation="MGQGRPRPSWSRWVCGRRRLGSRWAIPVVKLVGRRSLISLSSRV
                     VHGPFSEASQPVAAARGHRPHIHGLARERSDRAGRSPEKPVGRRSRRRSRRLCEAKSSE
                     YTQALSGPPEAPPCAAPVEPVGHQGGAGMAAYAIMRCKKLAKMGNVAASLKHAIRER
                     ETNPADARTPENEHWAASSTDEAMGRLELPEKRRKDAVLAVEYVMTASPEWWKSA
                     SQEQQAFFFEKAHKWLADKYGADRIVTASIHREDTSPHMTAFVVPITQDRLSAKEFI
                     GNKAQMTDRDQTTFAAAVADLGLQRGIEGSKARHTRIQAIFYEALERPPVGHVITISPAV
                     EPYAPYAPQGLAEKLGISKRVETPEAVADRLTKAVRGYEPALQAAAGAREMRKKADQA
                     QETARDLRELRKPLVDALGPLNRDMQAKAAAIKAVGEKLLTEQREVQRQKQAQRQQE
                     RGRAHFPEKCHLG*"
     promoter        2405..2454
                     /label="NEOKAN_promoter"
     gene            2546..3334
                     /gene="NeoR/KanR"
                     /label="NeoR/KanR"
     CDS             complement(2852..3388)
                     /translation="MAGWASLGRSFRTPESRSEELVKKAIEGDALRIGSGDTVKHEEA
                     VSPFAAKLFNSITGSQRYVLIARHTQPATVDESRKAAIFHHDIRQAGIAMGHDEILA
                     VGHARLEPGEQFGWREPLMLFVQIILIDKTGFHPSTCSLDAMFRLVVEWAGSRIRKMQ
                     PPHCISHDGYFLGRSKVR*"
                     /label="ORF frame 3"
     misc_feature    3499..3529
                     /note="Geneious type: operator"
                     /label="ArsR binding site"
     gene            3577..3930
                     /label="arsR"
     gene            3966..6059
                     /label="mtrB"

ORIGIN
1  gtgccggggc  cgttgagctt  cagcagcgag  atccagcgct  cggccaccaa  gtccttgact
61  gcgtatttga  ccgtccgcaa  agaacgtccg  atgagcttgg  aaagtgtctt  ctggctgacc
121  accacggcgt  tcgtggtgcc  catctgcgcc  acgaggtgat  gcagcagcat  tgccgcgcgt
181  gggttctcgt  caataaaccc  ggcccacgcc  tcatcgcttt  tgcgttccgt  ttgcacccag
241  tgaccgggct  tgttcttgcc  ttgaatcgcc  atttctctgg  actgcgtggc  catgcttctc
301  tccatcgcggt  aggggtcgcc  acggttgccg  caccatgcgc  aatcagctgc  aacttttctg
361  cagcgcgaca  acaattatgc  gttgcgtaaa  agtggcagtc  aattacagat  ttcttttaac
421  ctacgcaatg  agctatttgc  gggggtgccg  caatgagctg  ttgcgtatcc  ccttttttta
481  agttgttgat  ttttaagtct  ttgcatttgc  gccctatata  tagttctttg  gtgccccaa
541  aagggaaccc  ctgcgggggt  ccccccacgc  ttccgcgcgc  ctccccctcc  ggcaaaaaat
601  ggccccctcc  gggcttgggt  atcgactgcg  cggccttcgg  ccttgcccaa  ggtggcgctg
661  cccctcttga  acccccacgc  tcgcccgcgt  gaggctcggg  gggcaggcgg  gcgggcttcg
721  ccttcgactg  ccccactcgc  cataggcttg  ggtcgttcca  ggcgcgtcaa  ggccaagcgc
781  ctgcgcggtc  gctgcgcgag  ccttgaccgc  ccttccactt  ggtgtccaac  cggcaagcga
841  agcgcgccag  ccgcaggcgc  gaggcttttc  cccagagaaa  attaaaaaaa  ttgatggggc
901  aaggccgcag  ccgcgcagtc  tggagccggt  gggatatgtg  tcgaaggctg  ggtagccggt
961  gggcaatccc  tgtggtcaag  ctgctgggca  ggccagccct  gtccatcagc  ttgtccagca
1021  ggggtgtcca  cgggcccagc  gaagcgagcc  agccggtggc  cgctcgcggc  catcgctccac
1081  atatccacgc  gctggcaagg  gagcgagcgc  accgcgcagg  gcgaagccgc  gagagcaagc
1141  cgttagggcg  ccgcagccgc  cgtaggcggt  cagcactttg  cgaagcaaa  tctagttagt
1201  ataactcaag  attgagtggc  ccgcgggagg  caccgccttg  cgctgcccc  gtcgagccgc
1261  ttggacacca  aaagggaggg  gcaggcatgg  cggcctacgc  gatcatgcga  tgcaagaagc
1321  tggcgaaaa  gggcaacgtg  gcggccagtc  tcaagcacgc  ctaccgcgag  cgcgagacgc
1381  ccaacgctga  cgccagcagg  acgcccagga  acgagcactg  ggcggccagc  agcaccgatg
1441  aagcgatggg  ccgactgcgc  gagttgctgc  cagagaagcg  gcgaaggac  gctgtgttgg
1501  cggtcgagta  cgtcatgacg  gccagcccg  aatggtgaa  gtcggccagc  caagaacagc
1561  aggcgcgcgt  cttcgagaag  gcgcacaagt  ggctggcgga  caagtacggg  gcgcatcgca
1621  tcgtgacggc  cagcatccac  cgtgacgaaa  ccagcccgca  catgaccgcg  ttcgtggtgc
1681  cgctgacgca  ggacggcagg  ctgtcgccga  aggagttcat  cggcaacaaa  gcgcagatga
1741  ccgcgaccca  gaccacgttt  gcggctgctg  tggccgatct  agggctgcga  cggggcatcg
1801  agggcagcaa  ggcacgtcac  acgcgcattc  aggcgtttct  cgaaggccctg  gagcggccac
1861  cagtgggcca  cgtccaccat  agcccgaag  cggtcgagcc  acgcgcctat  gcaccgcagg
1921  gattgcccga  aaagctggga  atctcaaa  cgcgtgagac  gccggaagcc  gtcggccgac
1981  ggctgacaaa  agcggttcgc  caggggtatg  agcctgccct  acaggcccg  gcaggagcgc
2041  gtgagatgcg  caagaagcc  gatcaagccc  aagagacggc  ccgagacctt  cgggagcgcc
2101  tgaagcccg  ctgggacgcc  ctggggccgt  tgaatcgga  tatgcaggcc  aaggccgcgc
2161  cgatcatcaa  ggcggtgggc  gaaaagctgc  tgacggaaca  gcgggaagtc  cagcgccaga

```

2221 aacaggccca ggcgcagcag gaacgcgggc ggcacattt ccccgaaaa tgccacctgg
2281 gatgaatgtc agctactggg ctatctggac aagggaatac gcaagcgcaa agagaaagca
2341 ggtagcttgc agtgggctta catggcgata gctagactgg cgggttttat ggacagcaag
2401 cgaaccggaa ttgccagctg gggcgccctc tggttaaggtt gggaagccct gcaaaagtaa
2461 ctggatggct ttcttgccgc caaggatctg atggcgcaag ggaacaagat ttgatcaaga
2521 gacaggatga ggaatgttgc gcatgattga acaagatgga ttgcacgcag gttctccggc
2581 cgcttgggtg gagaggctat tcggctatga ctgggcacaa cagacaatcg gctgctctga
2641 tgccgcgctg ttccggctgt cagcgacagg ggcggcggtt ctttttgcga agaccgacct
2701 gtccggtgcc ctgaatgaac tacaggacga ggcagcgagg ctatcggtgg tgccacgcag
2761 gggcggttct tgccagcctg tgctcgacgt tgctactgaa cggggaaggg actggctgct
2821 attggcgcaa gtcccggggc aggatctcct gtcactcac ctgtctcctg ccgagaaagt
2881 atccatcatg gctgatgcaa tgcggcggtt gcatacgtt gatccggcta cctgccatt
2941 cgaccaccaa gcgaacatc gcatcgagcg agcacgtact cggatggaa gctgccatt
3001 cgatcaggat gatctggacg aagagcatca ggggctcgcg ccagccgaac tgttcgcca
3061 gctcaaggcg cgcatgccc acggcgagga tctcgtcgtg acccatggcg atgctgctt
3121 gccaatgttc atggtgaaa atggcgctt ttctggattc atcgactgtg cggcgctggg
3181 tgtggcgagc cgctatcagg acatagcggt ggctaccgtg gatattgctg aagagcttgg
3241 cggcgaaatg gctgaccgct tctcgtgct ttacggatc ggcgctcccg attcgacgag
3301 catcgccctc tatcgccctc ttgacgagtt cttctgagcg ggactctggg gttcgaagt
3361 accgaccaag cgacgcccac cctgcatca cgagatttgc attccaccgc cgccttctat
3421 gaaaggttgg actagagaat tcggcgccgc ttctagagcc aactcaaat tcacacctat
3481 taccttcttc tgcaactaca cattcgttaa gtcataatgt ttttgactt atccgcttcg
3541 aagagagaca ctacctgcaa caatcaggag cgcaatatgt catttctgtt acccatccaa
3601 ttgttcaaaa ttcttctgta tgaaccgct ctgggcatcg ttttactgct cagcgaaact
3661 ggagagttat gcgtctcgca tctctgcact gctctcgacc agtcgacgcc caagatctcc
3721 cgccacctgg cattgtctcg tgaagcggg ctattgctgg accgcaagca aggttaagtgg
3781 gttcattacc gcttatcacc gcatattcca gcatggcgcg cgaaaattat tgatgagggc
3841 tggcgatgtg aaacaggaaa gtttcaggcg attgtccgca accgtgctcg acaaaactgt
3901 tccggggaca gtaagaacat ttgcagttaa cggcgcgctc actagagaaa gaggagaaat
3961 actagatgaa atttaacctc aatttgatca ctctagcggtt attagccaac acaggtctgg
4021 ccgtcgtcgc tgatgggtat ggtctagcga atgccaatc tgaaaaagtg aaattatccg
4081 catggagctg taaaggctgc tgcgttgaac cgggcacatc aggcactgtg ggtgtcgggt
4141 tcggttaata cagcgaaagc gatattcgct ctgccaatgc ctttggtaca tccaatgaag
4201 tggcgggtaa atttgatgcc gatttaaaat ttaaagggtg aaagggttat cgtgccagt
4261 ttgatgctta tcaactcgtt atggatggcg gtcgcttaga tgcgaatgag ggcaaacagc
4321 gccagtacaa cgtcaatgtg aactatcgcc aaattgttac ctacgacagc aatagcgccc
4381 tatcgcccta cggcggtatt ggtggcaata acctcacgtt accgataaac tggataaacg
4441 caggttcaag caaccaaatg ccactcttga tggacagcct caatgccctc gaactctcac
4501 ttaaacgtga ggcgcagggg ttgggatttg aatatcaag tgaatccctg tggagcaact
4561 atgttaacta catcgctgaa gagaaaaccg gcttaaaaca agcctctggt agcttcttca
4621 accaatcgat tagtttagca tattacgaac gagccggtgg attacaccac tgacaccatt
4681 tcaaaactca ggtgtatcgt tgggttaccg cactcagtta caatgggtgca atattcaaaa
4741 acgaatacaa ccaattggac tttgaaaatg cttttaacc cactcttgggt gctcaaaccc
4801 aaggtacgat ggcactcgat ccggataaac agtcacacac cgtgtcgctg atgggacagt
4861 acaacgtagg cagcaacgca ctgtcgggtg gtattctgac cggacaaatg agccaaagtc
4921 aggcgttagt gacggataac taccgttatg ctaatcagct caataccgat cgcgtcgatg
4981 ccaaagtcga tctactgggt atgaacctga aagtctgtag caaagtgagc aatgatcttc
5041 gctttaacag tagttacgat tattacgaac gtgacaataa taccgaagta gaagaatgga
5101 ctacagatcag catcaacaat gtcaacggta aggtggctta taacaccctc tacgataatc
5161 gtacgcaacg cttttaaagt gccgcagatt atcgattac ccgcatatc aaactcgatg
5221 ttggttatga cttcaaacgt gaccaacgtg attatcaaga ccgtgaaacc acggatgaaa
5281 ataccgtttg ggcggcttta cgtgtaacaa gcttcgatac ttgggacatg tgggtaaaaa
5341 gcagttacgg taaccgtgac ggctcacaat accaagcgtc tgaatggacc tctctgaaa
5401 ccaacagcct gttacgtaag tacaatctgg ctgaccgtga cagaactcaa gtcgaagcac
5461 ggatcaccga ttccgcatga gaaagcctga ctatcgatgt tgggtcccggt tacgcgttag
5521 atgattatac cgatactgtg attggattaa ctgagtcaaa agacaccagt tatgatgcca
5581 acatcagtta tatgatcacc gctgacttac tggcaaccgc cttctacaat taccaaacca
5641 ttgagtctga acaggcggtt agcagcaatt acagcaccgc aacgtggaca ggttttatag
5701 aagatcaggt agatgtggtc ggtgcaggtt tcagctacaa caatctgctg gagaacaagt
5761 tacgcctagg actggactac acctattcca actccgacag taacactcaa gtcagacaag
5821 gtatcactgg cgactatggt gattattttg ccaaagtgcga taacattaac ttatacgtc
5881 aatatcaagc caccgagaaa ctgcgctgc gcttcgatta caaaattgag aactataagg
5941 acaatgacgc cgcaaatgat atgcgcgttg atggcatttg gaacgtcgta ggttttggt
6001 gtaacagcca tgactacacc gcacaaatgc tgatgctgag catgagttac aaactcta
6061 actagatgca cactgggtca ccttcgggtg ggcccttctg cgtttatata ctagagagag
6121 aatataaaaa gccagattat taatccggct tttttattat ttactagta ggcggcgctg
6181 cagtctagta tagtctggaa cagcgcaact acgggttgct gcgcaaccga agtgctaccg
6241 ggcggcgagc gtgaccgctg tcggcggtc caacggctcg ccatcgcca gaaaacacgg
6301 ctcatcgggc atcgcgagc gctgctgcc gcgcggttcc cattectcgt ttcgggtcaa
6361 ggtggtcagg tctggttcca tgcccggaat gccgggctg ctggcggtc cctcgccggg
6421 gccggctcgt agttgctgct cgcggcgata cagggtcggg atgcccgcga ggtcgccatg
6481 cccaacagc gatcgtcctt ggtcgtcgtg atcaaccacc acggcgccac tgaacaccga
6541 caggcgcaac tggtcgctgg gctggcccca gcgcacgcgg tcattgacca cgtaggccga
6601 cacc

//

6. REFERENCES

- Belkin, S., 2003. *Curr. Opin. Microbiol.* 6, 206–212.
- Bjerketorp, J., Håkansson, S., Belkin, S., Jansson, J.K., 2006. *Curr. Opin. Biotech.* 17, 43–49.
- Coursolle, D., Baron, D.B., Bond, D.R., Gralnick, J.A., 2010. *J. Bacteriol.* 192, 467–74.
- Coursolle, D., Gralnick, J.A., 2012. *Front. Microbio.* 3 (56).
- Dehio, C., Meyer, M., 1997. *J. Bacteriol.* 179, 538–40.
- Diorio, C., Cai, J., Marmor, J., Shinder, R., DuBow, M.S., 1995. *J. Bacteriol.* 177, 2050–2056.
- D'Souza, S.F., 2001. *Biosens. Bioelectron.* 16, 337–353.
- Friedman, E.S., Rosenbaum, M.A., Lee, A.W., Lipson, D.A., Land, B.R., Angenent, L.T., 2012a. *Biosens. Bioelectron.* 32, 309–13.
- Friedman, E.S., TerAvest, M.A., Venkataraman, A., Angenent, L.T., 2012b. Enzymatic and Microbial Electrochemical Systems, in Heldman, D.R. And Moraru, C.I. (Eds.), *Encyclopedia of Agricultural, Food, and Biological Engineering, Second Edition*. Taylor & Francis, New York.
- Goldbeck, C., Jensen, H., TerAvest, M.A., Beedle, N., Appling, Y., Helper, M., Cambray, G., Mutalik, V., Angenent, L.T., Ajo-Franklin, C.M., 2013. *ACS Synth. Biol.* 2, 150–159.
- Golitsch, F., Bücking, C., Gescher, J., 2013. *Biosens. Bioelectron.* 47, 285–291.
- Heitzer, A., Webb, O.F., Thonnard, J.E., Gary, S., Sayler, G.S., 1992. *Appl. Environ. Microb.* 58, 1839–1846.
- Joshi, N., Wang, X., Montgomery, L., Elfick, A., French, C.E., 2009. *Desalination.* 248, 517–523.
- Kim, B.H., Chang, I.S., Gil, G.C., Park, H.S., Kim, H.J., 2003. *Biotechnol. Lett.* 25, 541–545.
- Kim, H., Hyun, M., Chang, I., Kim, B., 1999. *J. Microbiol. Biotechnol.* 9, 354–367.
- King, J.M., Digrazia, P.M., Applegate, B., Burlage, R., Sanseverino, J., Dunbar, P., Larimer, F., Sayler, G.S., 1990. *Science.* 249, 778–81.
- Kovach, M.E., Elzer, P.H., Hill, D.S., Robertson, G.T., Farris, M.A., Roop, R.M., Peterson, K.M., 1995. *Gene.* 166, 175–176.
- Lei, Y., Chen, W., Mulchandani, A., 2006. *Anal. Chim. Acta.* 568, 200–210.
- McGuigan, C.F., Hamula, C.L.A., Huang, S., Gabos, S., Le, X.C., 2010. *Environ. Rev.* 18, 291–307.
- Myers, C.R., Nealson, K.H., 1988. *Science.* 240, 1319–1321.
- Ramanathan, S., Shi, W., Rosen, B.P., Daunert, S., 1997. *Anal. Chem.* 69, 3380–3384.
- Rosenbaum, M., Cotta, M.A., Angenent, L.T., 2010. *Biotechnol. Bioeng.* 105, 880–888.
- Sagi, E., Hever, N., Rosen, R., Bartolome, A.J., Rajan Premkumar, J., Ulber, R., Lev, O., Scheper, T., Belkin, S., 2003. *Sensor. Actuat. B-Chem.* 90, 2–8.
- Shin, H.J., 2011. *Appl. Microbiol. Biot.* 89, 867–877.
- Stookey, L.L., 1970. *Anal. Chem.* 42, 779–781.
- TerAvest, M.A., Li, Z., Angenent, L.T., 2011. *Energ. Environ. Sci.* 4, 4907–4916.
- Wu, N., Matand, K., Kebede, B., Acquaaah, G., Williams, S., 2010. *Electron. J. Biotechnol.* 13, 1–9.
- Vick, J.E., Johnson, E.T., Choudhary, S., Bloch, S.E., Lopez-Gallego, F., Srivastava, P., Tikh, I.B., Wawrzyn, G.T., Schmidt-Dannert, C., 2011. *Appl. Microbiol. Biotechnol.* 92, 1275–1286.